

Exclusive ϕ production in proton-proton collisions in the resonance model.

Amand Faessler^a, C. Fuchs^a, M.I. Krivoruchenko^b, B.V. Martemyanov^{a,b}.

^a *Institut für Theoretische Physik der Universität Tübingen, Auf der Morgenstelle 14, D-72076
Tübingen, Germany*

^b *Institute for Theoretical and Experimental Physics, B. Cheremushkinskaya 25, 117259 Moscow,
Russia*

Abstract

The exclusive ϕ meson production in proton-proton reactions is calculated within the resonance model. The considered model was already successfully applied to the description of π , η , ρ , ω , $\pi\pi$ production in proton-proton collisions. The only new parameter entering into the model is the $\omega - \phi$ mixing angle θ_{mix} which is taken equal to $\theta_{mix} \approx 3.7^\circ$.

I. INTRODUCTION

The production of ϕ mesons is strongly suppressed compared to that of ω mesons. This fact is known under the name “OZI rule” [1]. According to the OZI rule ϕ mesons can only be produced due to a small admixture of non-strange light quarks in their wave function. The corresponding mixing angle θ_{mix} is equal to $\theta_{mix} \approx 3.7^\circ$ [2]. According to this mixing angle the ratio of ϕ and ω mesons cross sections should at comparable energies naively be equal to

$$R_{\phi/\omega} = \tan^2 \theta_{mix} \cdot F \approx 4.2 \times 10^{-3} \cdot F , \quad (1)$$

where F is a correction due to the different phase space factors for ϕ and ω mesons. Experimentally, the ratio $R_{\phi/\omega}$ is in $pp \rightarrow pp\phi(\omega)$ reactions, however, known to be one order of magnitude larger than the naive expectation [3–7]. Various models have been used as a framework for theoretical attempts to explain this large discrepancy, ranging from simple one-pion exchange models [8] to more sophisticated effective Lagrangian models [9]. Also the resonance model is successfully explaining the production of a large variety of mesons such as π , η , ρ , ω , $\pi\pi$ in elementary nucleon-nucleon reactions [10,11]. In the present note we extend the application of the resonance model to the description of ϕ meson production in pp collisions. In the next section we briefly describe the resonance model and present the results for the ϕ meson production. Then we discuss the results and conclude.

II. RESONANCE MODEL

The production of ϕ mesons is described by a two step mechanism, i.e. the excitation of nucleon resonances and their subsequent decay $pp \rightarrow pR \rightarrow pp\phi$. The cross sections for the resonance production were taken from ref. [10] where they were fitted to describe π , η , ρ , $\pi\pi$ production in NN collisions. The ϕNR coupling is obtained from the known ωNR coupling of the ω meson [11] and the mixing angle between ϕ and ω mesons. The ωNR couplings, in turn, have been determined within the framework of the extended Vector Meson

Dominance (eVMD) model by fitting the available data on electro- and photo- production of nucleon resonances as well as their mesonic decays [12]. The description of the ϕ and ω meson production in elementary nucleon-nucleon reactions is then essentially parameter free since all model parameters have already been fixed by other sources. In [11] it was demonstrated that available data on the exclusive ω production in pp reactions are very accurately reproduced by the present model over a wide energy range, i.e. from extremely close to threshold up to several GeV above threshold.

The $pp \rightarrow pR \rightarrow pp\phi$ cross section is given as follows:

$$\sigma(s)^{pp \rightarrow pp\phi} = \sum_R \int_0^{(\sqrt{s}-2m_p)^2} dM^2 \int_{(m_p+M)^2}^{(\sqrt{s}-m_p)^2} d\mu^2 \frac{d\sigma(s, \mu)^{pp \rightarrow pR}}{d\mu^2} \frac{dB(\mu, M)^{R \rightarrow p\phi}}{dM^2} , \quad (2)$$

with M being the running mass of ϕ meson. The cross sections for the baryon resonances production are given by

$$d\sigma(s, \mu)^{pp \rightarrow pR} = \frac{|\mathcal{M}_R|^2}{16p_i \sqrt{s} \pi^2} \Phi_2(\sqrt{s}, \mu, m_p) dW_R(\mu) \quad (3)$$

with $\Phi_2(\sqrt{s}, \mu, m_p) = \pi p^*(\sqrt{s}, \mu, m_p)/\sqrt{s}$ being the two-body phase space, $p^*(\sqrt{s}, \mu, m_p)$ the final c.m. momentum, p_i the initial c.m. momentum. μ and m_R are the running and pole masses of the resonances, respectively, m_p the proton mass. The mass distribution $dW_R(\mu)$ of the resonances is described by the standard Breit-Wigner formula:

$$dW_R(\mu) = \frac{1}{\pi} \frac{\mu \Gamma_R(\mu) d\mu^2}{(\mu^2 - m_R^2)^2 + (\mu \Gamma_R(\mu))^2} . \quad (4)$$

The sum in (2) runs over the same set of nucleon resonances which is responsible for the ω meson production [11]. This includes all well established (4*) N^* resonances quoted by the PDG [2] with masses below 2 GeV. The branching to the ϕ decay mode is given by

$$dB(\mu, M)^{R \rightarrow p\phi} = \frac{\tan^2 \theta_{mix} \Gamma_{N\omega}^R(\mu, M)}{\Gamma_R(\mu)} dW_\phi(M) , \quad (5)$$

with $\Gamma_{N\omega}^R(\mu, M)$ calculated the same way as in the case of ω meson production [11].

The ϕ mass distribution $dW_\phi(M)$ is also described by a Breit-Wigner distribution, i.e. substituting $R \rightarrow \phi$ and $\mu \rightarrow M$ in Eq. (4). The energy dependence of the ϕ width $\Gamma_\phi(M)$ can be calculated according to

$$\Gamma_\phi(M) = \Gamma_\phi(m_\phi) \left(\frac{p^*(M, m_K, m_K)}{p^*(m_\phi, m_K, m_K)} \right)^3 \left(\frac{m_\phi}{M} \right)^2. \quad (6)$$

Important properties of the $\Gamma_{N\omega}^R(\mu, M)$ width are the following ones: the M dependence of magnetic, electric and Coulomb couplings entering into the amplitude and the Blatt-Weisskopf suppression factor which suppresses the width for large off-shell masses μ of the resonances [11]. Their combined effects lead to an increase of the $\Gamma_{N\omega}^R(\mu, M)$ width with M increasing from m_ω to m_ϕ for $\mu > 2GeV$. This effect is finally responsible for the violation of the naive OZI rule estimate (1).

The results for cross section $\sigma(pp \rightarrow pp\phi)$ are presented in Fig. 1 in form of the ratio of the ϕ over ω meson production cross sections. The ω mesons production cross section was shown to agree well with experimental data if one uses a strong $N^*(1535)N\omega$ coupling [11]¹. The dashed line in Fig. 1 corresponds to the idealized case of stable ϕ and ω mesons, i.e. to the limit of zero widths.

III. SUMMARY

We have calculated the cross section for ϕ meson production in pp collision within the resonance model. The experimental data are well reproduced. The large violation of the OZI estimate for the ϕ over ω production, observed experimentally, is explained without introducing additional parameters into the present model. A further test of the resonance model could be provided by a comparison of μ^2 distributions (the distribution of the invariant mass of proton- ϕ meson system) as soon as experimental data are available for this observable.

One of us (B.V.M.) is indebted to the Institute for Theoretical Physics of the University

¹Here we do not subtract contributions from off-shell ω production from the total cross section since at the considered energies, i.e. above the ϕ meson production threshold, these contributions are less than 10% of the total ω cross section. Hence the effect that can be approximatly neglected.

of Tübingen for kind hospitality. This work is supported by RFBR grant No. 03-02-04004, DFG grant No. 436 RUS 113/721/0-1.

REFERENCES

- [1] S. Okubo, Phys. Lett. **B5**, 165 (1963);
G. Zweig, CERN Report No.8419/TH, 412 (1964);
I. Iizuka, Prog. Theor. Phys. Suppl. **37 & 38**, 21 (1966).
- [2] Particle Data Group, Phys. Rev. D **66**, (2002); Eur. Phys. J. **C3**, 1 (1998).
- [3] DISTO Collaboration, F. Balestra et al., Phys. Rev. C **63**, 024004 (2001).
- [4] V. Blobel et al., Phys. Lett. **B59**, 88 (1975).
- [5] R. Baldi et al., Phys. Lett. **B68**, 381 (1977).
- [6] M.W. Arenton et al., Phys. Rev. D **25**, 2241 (1982).
- [7] S.V. Golovkin et al., Z. Phys. A **359**, 435 (1997).
- [8] A. Sibirtsev, Nucl. Phys. **A604**, 455 (1996).
- [9] K. Tsushima and K. Nakayama, nucl-th/0304017.
- [10] S. Teis, W. Cassing, M. Effenberger, A. Hombach, U. Mosel, and Gy. Wolf, Z. Phys. **A356**, 421 (1997).
- [11] C. Fuchs, M.I. Krivoruchenko, H.L. Yadav, Amand Faessler, B.V. Martemyanov and K. Shekhter., Phys. Rev. C **67**, 025202 (2003).
- [12] M. I. Krivoruchenko, B. V. Martemyanov, A. Faessler and C. Fuchs, Annals Phys. (N.Y.) **296**, 299 (2002).




FIG. 1. The ratio of the $\sigma(pp \rightarrow pp\phi)$ over $\sigma(pp \rightarrow pp\omega)$ cross sections as a function of the c.m. energy above the threshold $Q = \sqrt{s} - 2m_p - m_\phi$ (solid line) is compared to experimental data [3–6]. The dashed line corresponds to idealized case of stable ϕ and ω mesons, i.e. the zero width limit.